



Stationary states and dynamics of superconducting thin films

Ögren, Magnus; Sørensen, Mads Peter; Pedersen, Niels Falsig

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Ögren, M., Sørensen, M. P., & Pedersen, N. F. (2011). *Stationary states and dynamics of superconducting thin films*. Poster session presented at 7.th International Conference on Vortex Matter in Nanostructured Superconductors, Rhodes Greece,.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Stationary states and dynamics of superconducting thin films

A poster presentation at:

*7.th International Conference on Vortex Matter in Nanostructured Superconductors,
Rhodes Greece, 2011*

M. Ögren, M. P. Sørensen and N. F. Pedersen

Dep. of Mathematics, Technical University of Denmark, 2800 Kongens Lyngby, Denmark

Abstract

The Ginzburg-Landau (GL) theory is a celebrated tool for theoretical modelling of superconductors [1]. We elaborate on different partial differential equations (PDEs) and boundary conditions for GL theory, formulated within the finite element method (FEM) [2]. Examples of PDEs for the calculation of stationary states with the GL equation and with the time-dependent GL equation are given. Moreover we study real time evolution with the so called Schrödinger-GL equation [3]. For simplicity we here present numerical data for a two-dimensional rectangular geometry, but we emphasize that our FEM formulation can handle complex geometries also in a three-dimensional superconducting structure. To include external currents in our modelling we discuss the role of the boundary conditions for the external magnetic field [4]. Finally we show results for the pinning of vortices with controlled impurities.

References

- [1] V. L. Ginzburg and L. D. Landau, Zh. Eksp. Teor. Fiz. **20**, 1064 (1950).
- [2] T. S. Alstrøm *et al.*, Acta Appl. Math., Springer Netherlands (2010).
- [3] L. Peres and J. Rubinstein, Physica D **64**, 299 (1993).
- [4] M. Machida and H. Kaburaki, Phys. Rev. Lett. **71**, 3206 (1993).